




BMJ Open Ketamine for the treatment of prehospital acute pain: a systematic review of benefit and harm

Mårten Sandberg,^{1,2} Per Kristian Hyldmo,^{3,4,5} Poul Kongstad,⁶ Kristian Dahl Friesgaard,^{7,8} Lasse Raatiniemi ,^{9,10} Robert Larsen,^{11,12} Vidar Magnusson,¹³ Leif Rognås ,^{14,15,16} Jouni Kurola,^{17,18} Marius Rehn ,^{1,3,4} Gunn Elisabeth Vist¹⁹

To cite: Sandberg M, Hyldmo PK, Kongstad P, *et al.* Ketamine for the treatment of prehospital acute pain: a systematic review of benefit and harm. *BMJ Open* 2020;**10**:e038134. doi:10.1136/bmjopen-2020-038134

► Prepublication history and additional material for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-038134>).

Received 28 February 2020
Revised 19 September 2020
Accepted 30 October 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Marius Rehn;
rehmar@snla.no

ABSTRACT

Background Few publications have addressed prehospital use of ketamine in analgesic doses. We aimed to assess the effect and safety profile of ketamine compared with other analgesic drugs (or no drug) in adult prehospital patients with acute pain.

Methods A systematic review of clinical trials assessing prehospital administration of ketamine in analgesic doses compared with other analgesic drugs or no analgesic treatment in adults. We searched PubMed, EMBASE, Cochrane Library and Epistemonikos from inception until 15 February 2020, including relevant articles in English and Nordic languages. We used the Cochrane and Grading of Recommendations Assessment, Development and Evaluation methodologies and exclusively assessed patient-centred outcomes. Two independent authors screened trials for eligibility, extracted data and assessed risk of bias.

Results We included eight studies (2760 patients). Ketamine was compared with various opioids given alone, and intranasal ketamine given with nitrous oxide was compared with nitrous oxide given alone. Four randomised controlled trials (RCTs) and one cluster randomised trial included 699 patients. One prospective cohort included 27 patients and two retrospective cohorts included 2034 patients. Five of the eight studies had high risks of bias. Pain score with ketamine is probably lower than after opioids as demonstrated in a cluster-RCT (308 patients) and a retrospective cohort (158 patients) study, Δ visual analogue scale -0.4 (-0.8 to 0.0) and Δ numeric pain rating scale -3.0 (-3.86 to -2.14), respectively. Ketamine probably leads to less nausea and vomiting (risk ratio (RR) 0.24 (0.11 to 0.52)) but more agitation (RR 7.81 (1.85 to 33)) than opioids.

Conclusions This systematic literature review finds that ketamine probably reduces pain more than opioids and with less nausea and vomiting but higher risk of agitation. Risk of bias in included studies is high.

Other Scandinavian society of anaesthesiology and intensive care medicine funded meetings and software. The Norwegian Air Ambulance Foundation funded publication. Otherwise this research received no grant from any agency in the public, commercial or not-for-profit sectors.

PROSPERO registration number CRD42018114399.

Strengths and limitations of this study

- A systematic review where main outcomes were assessed according to the Grading of Recommendations Assessment, Development and Evaluation method.
- Studies were heterogeneous in terms of setting, patient population, outcomes and comparators.
- Only English-language and Scandinavian-language articles were included.

INTRODUCTION

Prehospital acute pain is a frequent symptom and often inadequately managed.^{1–3} Several analgesics are administered by prehospital emergency medical services throughout the world without solid evidence of their efficacy and safety. The heterogeneity in pain management strategies may reflect the varying competence levels of providers ranging from technicians with basic training to specially trained physicians. Opioids are most frequently used, but their cerebral, haemodynamic and respiratory side effects remain a potential challenge in unstable and undifferentiated prehospital patients.⁴ Ketamine is an alternative to opioids. Ketamine exerts its effects mainly as an N-methyl-D-aspartate antagonist and, depending on the dose, can be considered as an analgesic, a sedative or an anaesthetic drug.⁵ One attractive feature for prehospital use of ketamine is its ability to preserve upper airway reflexes. Respiratory rate may increase, and ketamine can cause bronchodilation. While ketamine generally preserves respiratory function, ketamine can cause respiratory depression if given quickly.⁶ There is a risk of laryngospasm, which may require intubation in a very small fraction of cases.⁷

Ketamine can be administered in a variety of routes, most commonly intramuscularly,

intranasally and intravenously, although per oral and per rectal doses are used in different settings. Although originally believed to cause an increase in intracranial pressure (ICP), recent work in critical care patients indicates that ketamine has little or no impact on ICP. In two studies comparing ketamine and sufentanil, the authors concluded that ketamine did not affect ICP and that it was safe to administer to patients with traumatic brain injury (TBI).^{8 9} In another study, ketamine in conjunction with propofol was administered to patients with TBI, and a significant decrease in ICP was recorded.¹⁰ In one study on children with TBI, a reduction in ICP by up to 30% was found, and cerebral perfusion was improved.¹¹ In these studies, ketamine was used in anaesthetic doses, and the results should be interpreted with caution.

Moderate or severe agitation occurs in 5%–30% of adult patients; some clinicians administer boluses of midazolam to avoid this phenomenon.⁶ A randomised controlled trial (RCT) showed that this practice significantly reduced agitation in adults; however, one trial found that it did not reduce agitation in children.^{6 12}

Proper pain relief allows prehospital care providers to meet essential clinical endpoints, for example, facilitating fracture manipulation. Although analgesia should be titrated for the desired effect, pain relief is frequently suboptimal, possibly due to concerns about adverse events.¹³ Ketamine may be a useful prehospital analgesic mainly due to its ability to provide excellent analgesic effects with a lower incidence of respiratory depression than that caused by opioids. These positive effects have been demonstrated in fracture management,¹⁴ burn treatment¹⁵ and traumatic amputation.¹⁶

The aim of this systematic literature review is to explore the benefit and harm of ketamine compared with other analgesic drugs (or no drug) in prehospital patients with acute pain.

METHODS

We conducted this systematic review according to the Cochrane Handbook for systematic reviews of interventions¹⁷ and as described in our protocol as specified below.

Inclusion criteria

We used the following inclusion criteria:

Population

Adult patients (18 years of age or older) with acute pain in the prehospital setting

Intervention

Ketamine

Comparison

Other analgesics, no analgesics or ketamine given in another dose or another route of administration or ketamine given in combination with other analgesics

Outcomes

Pain reduction, speed of onset, duration of effect and relevant adverse events such as mortality, morbidity, anaphylaxis, nausea and vomiting, hypotension, respiratory failure, loss of airway patency, emergence phenomena (as defined by study authors).

We included all adult patients (18 years of age or older) with acute pain, regardless of aetiology, managed in the prehospital setting. We also sought to identify adverse effects that are not previously reported. The following study designs were considered eligible for inclusion in the meta-analysis: RCTs, non-randomised controlled studies, cohort studies with a control group, interrupted time series and controlled before-and-after studies. Case series were also included for information relating to safety. Systematic reviews of high-quality answering to our inclusion criteria were evaluated for eligible studies. Other systematic reviews would have been used to check for relevant references.

Exclusion criteria

Children (younger than 18 years of age) and patients with chronic pain and/or patients who used ketamine as part of their regular treatments were not included in this review. We excluded all studies that were not conducted in the prehospital setting as well as conference abstracts, letters and publications without full texts available.

Search strategy

An experienced research librarian in collaboration with the authors developed the search strategy based on the inclusion criteria. The following databases were searched from their inception: PubMed, EMBASE, Cochrane Library and Epistemonikos.

The most recent search was conducted on 15 February 2020, and the full search strategy is presented in online supplemental appendix 1. The search was limited to the following languages: Danish, English, Finnish, Icelandic, Norwegian and Swedish.

The reference lists of the included publications were checked in order to identify relevant articles not found in the original search.

Study selection

For each step in the review process, no assessor handled publications they had coauthored. MS and either PKH, MR or PK independently assessed all titles and abstracts identified from the search according to the inclusion criteria above. References that were considered potentially relevant were collected, and the full-text articles were assessed independently by two assessors using the same inclusion criteria. Any disagreement between the initial two assessors was discussed and resolved by all assessors. The process of study selection based on titles and abstracts, study selection based on full-text articles and risk of bias assessments were conducted using Covidence (Covidence systematic review software, Veritas Health

Innovation, Melbourne, Australia. Available at www.covidence.org).

Assessment of risk of bias

MS and either PKH, MR or PK independently assessed the risk of bias for each of the included studies in accordance with the recommendations by the Cochrane Collaboration.¹⁷ For RCTs, the following items were assessed for risk of bias: (1) sequence generation, (2) concealment of allocation, (3) blinding of participants and personnel, (4) blinding of outcome assessor, (5) incomplete outcome data, (6) selective outcome reporting and (7) other risk of bias. For non-RCTs and other studies with a control group, the following items were also assessed for risk of bias: (8) similarity of baseline characteristics, (9) similarity of baseline outcome data and (10) contamination. All items were rated as either high, unclear or low risk of bias.

Data extraction

MS and either PKH or PK independently extracted data from each included study. We extracted data pertaining to full references; study design and country in which the study was conducted; characteristics of the population, for example, number of patients; age; gender; cause of pain; setting and context; type and dose of analgesics

given; cadre/competency of the healthcare personnel who administered the analgesic; comparison/control intervention; attrition; outcomes and follow-up times. We did not contact any study investigators to obtain information not described in the original articles.

The process of study selection based on titles and abstracts, study selection based on full-text articles as well as risk of bias assessments were conducted using Covidence.

Statistical analysis

Dichotomous outcomes are presented as risk ratios (RRs) with 95% CIs. Continuous outcomes are presented as the mean difference between the groups with 95% CIs. If different scales were used to measure the same outcome, we would have calculated standardised mean difference with a 95% CI. We used Review Manager (RevMan V.5.3) software to generate forest plots. Attrition was handled using intention-to-treat analysis. We evaluated statistical heterogeneity using the Q test and I^2 -statistic. Analysis was by inverse variance and random effects methods. Zero events were presented descriptively.

Grading our confidence in the evidence

We assessed our confidence in the evidence for each outcome using the Grading of Recommendations




Figure 1 Flow diagram of evaluated records.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias	Baseline similarities
Randomised controlled trials								
Andolfatto 2019	+	+	+	?	+	+	+	+
Galinski 2007	+	+	?	?	+	+	?	+
Jennings 2012	?	+	-	-	+	+	+	+
Tran 2014	-	-	-	-	+	+	+	+
Wiel 2015	+	+	+	-	+	?	+	+
Cohorts								
Bronsky 2019	NA	NA	+	-	+	+	+	+
Johansson 2009	NA	NA	-	-	+	+	+	+
Losvik 2015	NA	NA	+	-	+	?	+	-

Figure 2 Risk of bias.

Assessment, Development and Evaluation (GRADE) method.¹⁸ Our confidence is presented as high, moderate, low or very low. The evidence across each outcome is assessed by eight criteria. Five criteria lowered our confidence in the evidence: (1) risk of bias/methodological limitations, (2) consistency between studies (statistical heterogeneity), (3) directness (similar study participants, intervention, comparator and outcome measures in the included studies to the population and target interventions and measures), (4) precision of results and (5) reporting bias. Three criteria were used to consider upgrading evidence from observational studies that had not been downgraded: (1) strong or very strong association between intervention and outcome;; (2) large or very large dose response and (3) situations where all plausible confounders would have reduced the effect. For questions about the effect of interventions, RCTs started at high confidence, and observational studies started at low confidence.

Breach of protocol

We did make a breach of protocol; the largest study (Losvik *et al*) we included also contained treatment data from a few children.¹⁹

Patient and public involvement

The development of the research question and outcome measures were informed by studies indicating that prehospital acute pain is a frequent symptom and often

inadequately managed.¹⁻³ No patients were directly involved in the design or conduct of this study.

The results will be disseminated as a part of a Scandinavian society of anaesthesiology and intensive care medicine guideline on prehospital pain management.

RESULTS

The systematic literature search identified 1197 references; we considered 60 to be potentially relevant and assessed those publications in full. We included seven of these studies in the final analysis. In addition, two unique references in the reference lists of the seven publications were assessed and one of the references was also included. **Figure 1** shows the flow diagram of the identified references. The 53 studies that were assessed in full-text articles and excluded are presented in online supplemental appendix 2 with the reason for their exclusion.

Characterisation of the trials

The eight included studies were conducted in Australia,²⁰ Canada,²¹ France,²²⁻²³ Iraq,¹⁹ Sweden,²⁴ USA²⁵ and Vietnam.²⁶ A total of 2760 prehospital patients with acute pain were included in these eight studies. Four RCTs²⁰⁻²³ and one cluster randomised trial²⁶ included 699 patients. One prospective cohort study²⁴ included 27 patients. Two retrospective cohort studies¹⁹⁻²⁵ included 2034 patients. The largest of these studies, with 1876 patients, was conducted in the war zones and mine fields of northern and central Iraq.¹⁹ Two authors stated that their studies were conducted in rural areas, with one in Australia and one in Vietnam. The latter study included areas with mine fields, and three patients had been involved in mine accidents. This study also included children; however, the vast majority of included patients was probably adults because the mean ages of the groups were 35.5 years and 36.9 years. Therefore, this study was included. The reported time frame was similar in all studies; that is, from drug administration to admission to hospital.

Risk of bias assessment

Our assessments regarding each bias domain are provided in **figure 2**. Three of the five RCTs had a high risk of bias, with the main reasons being lack of random sequence generation, lack of allocation concealment or lack of blinding of patients, personnel and outcome assessors.

Comparisons

The included studies covered five comparisons involving ketamine (**table 1**):

- Ketamine intravenous versus opioids (morphine,²⁶ fentanyl,²⁵ pentazocine¹⁹) intravenous.
- Ketamine intravenous and morphine intravenous versus only morphine intravenous.²⁰⁻²³⁻²⁴
- Ketamine intravenous given as continuous administration versus ketamine intravenous as single dose.²²
- Ketamine intranasally and nitrous oxide versus only nitrous oxide.²¹

Table 1 Summary of included studies

Reference Study design Country	Ketamine	Comparison	Outcomes
Ketamine intravenous vs opioids intravenous			
Bronsky <i>et al</i> ²⁵ 2019 Retrospective cohort USA	n=79, ketamine 0.3 mg/kg intravenous every 20 min as needed, maximum three doses	n=79, fentanyl 2 µg/kg bolus intravenous over 1 to 2 min with additional dose every 10 min as needed	Change in pain scores, serious adverse events, GCS
Losvik <i>et al</i> ¹⁹ 2015 Retrospective cohort Iraq	n=713, ketamine 0.2 mg/kg intravenous, in case of unrest, 5 mg diazepam intravenous. During protracted evacuations with repeated ketamine doses, 1 mg atropine was administered. Repeat doses of ketamine allowed.	n=888, pentazocine 0.4 mg/kg intravenous for adults, repeat doses allowed	Change in physiological severity score
Tran <i>et al</i> ²⁶ 2014 Cluster—RCT Vietnam	n=169, ketamine 0.2 to 0.3 mg/kg was administered as slow intermittent intravenous injections	n=139, morphine administered in one single i.m. dose; 10 mg for adult patients, 5 mg for paediatric patients	Change in pain score, serious adverse events, adverse events, satisfaction, mean treatment time (head trauma)
Ketamine and morphine intravenous vs morphine intravenous alone			
Galinski <i>et al</i> ²³ 2007 RCT France	n=33, ketamine 0.2 mg/kg intravenous in 3 mg morphine every 5 min if necessary	n=32, morphine 3 mg intravenous every 5 min if necessary	Change in pain score, adverse events
Jennings <i>et al</i> ²⁰ 2012 RCT Australia	n=70, morphine 5 mg intravenous initial dose followed by a ketamine bolus of 10 or 20 mg according to body size, followed by 10 mg ketamine every 3 min thereafter until pain was relieved	n=65, morphine 5 mg intravenous initial dose followed by 5 mg intravenous every 5 min until pain was relieved	Change in pain score, adverse events, GCS
Johansson <i>et al</i> ²⁴ 2009 Prospective cohort Sweden	n=16, morphine 0.1 mg/kg intravenous followed by ketamine 0.2 mg/kg if pain score ≥4 after 5 min	n=11, mg/kg morphine 0.1 mg/kg intravenous followed by morphine 0.1 mg/kg if pain score ≥4 after 5 min	Change in pain score, adverse event, mean treatment time
Ketamine continuous intravenous administration vs ketamine intravenous one dose			
Wiel <i>et al</i> ²² 2015 RCT France	n=30, all patients received ketamine 0.2 mg/kg intravenous bolus combined with morphine 0.1 mg/kg intravenous followed by ketamine 0.2 mg/kg/h. Additional morphine 0.05 mg/kg was allowed every 5 min if VAS>3/10	n=33, all patients received a ketamine 0.2 mg/kg intravenous bolus combined with morphine 0.1 mg/kg intravenous followed by a saline infusion of the same volume. Additional morphine 0.05 mg/kg was allowed every 5 min if the VAS>3/10	Change in pain score, adverse events, satisfaction
Intranasal ketamine and inhaled nitrous oxide vs only inhaled nitrous oxide			
Andolfatto <i>et al</i> ²¹ 2019 RCT Canada	n=60, all patients received approximately 0.75 mg/kg intranasal ketamine (30 mg for patients<50 kg, 50 mg for patients 50–100 kg, 75 mg for patients>100 kg) combined with inhaled nitrous oxide	n=60, all patients received inhaled nitrous oxide	Change in pain score, adverse events, satisfaction
Ketamine intravenous vs no analgesic treatment			
Losvik <i>et al</i> ¹⁹ 2015 Retrospective cohort Iraq	n=713, ketamine 0.2 mg/kg intravenous, in case of unrest, 5 mg diazepam intravenous. During protracted evacuations with repeated ketamine doses, 1 mg atropine was administered. Repeat doses of ketamine allowed.	n=275, no analgesic treatment	Change in physiological severity score

GCS, Glasgow coma scale; VAS, visual analogue scale.




Figure 3 Ketamine versus opioids—change in pain score.

► Ketamine intravenous versus no analgesia/no medication¹⁹

In table 1, we give a short description of the included studies and the doses used, while the excluded studies are presented with the reason for their exclusion in online supplemental appendix 2. One study contributed to two comparisons,¹⁹ meaning that 713 patients who received ketamine are compared two times, first with patients who received opioids and again with patients who did not receive analgesic treatment.

Ketamine versus opioids

A change in pain score was reported in two studies. Bronsky *et al*²⁵ used the numeric pain rating scale (NRS), where 1 represents no pain and 10 represents extreme pain, while Tran *et al*²⁶ measured the change in pain using the visual analogue scale (VAS) but did not explicitly give a range. Figure 3 shows that both studies reported a greater reduction in pain scores with ketamine than with the opioids fentanyl (Mean Difference (MD) -3.0 (95% CI -3.86 to 2.14)) and morphine (MD -0.4 (95% CI -0.08 to 0.0)).

The main outcome in the study by Losvik *et al*¹⁹ was the physiological severity score (PSS). The PSS was calculated from the blood pressure, respiratory rate and consciousness level.²⁷ They reported exactly the same change, at 1.5 (95% CI 1.4 to 1.6), in the PSS for both the ketamine and the pentazocine groups. Hence, no difference was found between the treatment groups.

Adverse events were reported in the Vietnamese study²⁶; fewer patients with nausea and vomiting were found in the ketamine group than in the morphine group and fewer patients with agitation were found in the morphine group than in the ketamine group (figure 4).




Figure 4 Ketamine versus opioids—adverse events.

In the study where ketamine and fentanyl were compared,²⁵ four adverse events were reported: two patients experienced respiratory compromise and two patients suffered haemodynamic instability. All four patients were in the fentanyl group.

The change in Glasgow coma scale (GCS) was measured by Bronsky *et al*²⁵ and found to be similar for ketamine and fentanyl, MD -0.13 (95% CI -0.33 to 0.07).

Ketamine and morphine versus only morphine

Changes in the pain scores were measured in both the Australian²⁰ and in the Swedish study²⁴ using a scale from 1 to 10 where 10 represented extreme pain. In the French study,²³ a scale from 0 to 100 was used, and we have transferred this to a 0 to 10 scale in order to include this study in the meta-analysis. Figure 5 shows the change in the pain score when prehospital patients received both ketamine and morphine compared with patients who received only morphine. Although the RCT performed by Jennings *et al*²⁰ found lower pain scores in patients receiving combined ketamine and morphine than in patients receiving only morphine. When combined with the RCT by Galinski *et al*,²³ the meta-analysis shows a non-significant reduction (MD -1.51 (95% CI -3.36 to 0.33)) in pain score. The small prospective cohort²⁸ also found a non-significant reduction (MD -1.30 (95% CI -2.95 to 0.35)) in pain score.

Adverse events were measured in both studies, and the results are illustrated in figure 6. It is important to note that the nausea and vomiting are included in the total adverse events in the RCTs. These results are characterised by few events but indicate that morphine alone may lead to fewer adverse events than the combination of ketamine and morphine.




Figure 5 Combined ketamine and morphine compared with only morphine—change in pain score.

The RCT by Jennings *et al*²⁰ measured the GCS score and found that the median score was unchanged between initial assessment and the follow-up time, with a median score of 15 for both groups.

The French RCT reported use of fewer boluses of morphine when combined with ketamine (1 bolus (95% CI 0 to 2) compared with 2.3 boluses (95% CI 1.8 to 3.8) when using morphine alone).²²

The Swedish, prospective cohort by Johansson *et al*²⁸ reported a non-significant trend for shorter treatment time with morphine alone than with ketamine and morphine combined (10 min (95% CI -1.4 to 21.4)). Ketamine was administered nasally, thereby avoiding the need for intravenous access.

Continuous ketamine administration versus ketamine given as a bolus

One multicentre RCT conducted in France compared the continuous administration of ketamine with a bolus dose of ketamine but both groups also received morphine.²² Changes in pain were measured using a VAS from 0 to 10 (worst) and were similar in both groups (VAS -0.6 (95% CI -1.84 to 0.64)).

The main outcome of this study was the amount of additional morphine used (mg/kg) ($p=0.18$), indicating that there was no difference between the continuous group, at 0.048 (first quartile, third quartile 0.000, 0.150), and the bolus group, at 0.107 (first quartile, third quartile 0.052, 0.150). The duration of care for both groups was 35 min. Nausea and vomiting were not reported in patients in the

continuous group but were reported in three patients in the bolus group.

Ketamine and nitrous oxide versus only nitrous oxide

Andolfatto *et al* used a verbal NRS pain score and evaluated the scores after 15 min and 30 min.²¹ More patients in the ketamine and nitrous oxide group had a reduction in pain of 2 or more points than those in the saline and nitrous oxide group at both time points (figure 7).

They reported no serious adverse effects in either group, but a considerable number of minor adverse events, such as feeling of unreality, dizziness, nausea, fatigue, general discomfort, mood change, hallucination, change in hearing and headache, occurred. Most of these side effects (52 of 66 events) were reported in the group of patients who received ketamine and nitrous oxide combined, as shown in figure 8.

Ketamine versus no analgesic treatment

The retrospectively matched observational study of patients/causalities in the war zone in Iraq compared the use of ketamine with no analgesic treatment.¹⁹ The main outcome in this study was the PSS, which was calculated from the blood pressure, respiratory rate and consciousness level. There was a non-significant trend for lower PSS with ketamine compared with no analgesics (MD -0.2 (95% CI -0.42 to 0.02)).




Figure 6 Combined ketamine and morphine compared with only morphine—adverse events.




Figure 7 Ketamine and N2O versus only N2O—change in pain score.

Adverse events from ketamine across the five comparisons

Seven of the eight studies reported on adverse events and/or side effects from use of ketamine. Five studies reported on nausea and vomiting from ketamine alone (8/169),²⁶ from ketamine continuous administration (0/30),²² from ketamine bolus administration (3/33),²² and from combined ketamine and morphine (8/33),²³ (4/70),²⁰ (7/16).²⁴ Time for administering each bolus was not reported. Only one study reported on agitation, from ketamine alone (19/169).²⁶ Four studies reported adverse events, and two stated that nausea and vomiting were included as adverse events, from ketamine and morphine (21/33),²³ (27/70).²⁰ One study reported adverse events from ketamine and nitrous oxide (52/60)²¹ and one study reported no adverse events from ketamine alone (0/79).²⁵

GRADE

The quality of the main outcomes for the comparisons involving the use of ketamine for the treatment of prehospital acute pain was assessed according to the GRADE method.¹⁸ The quality of evidence could be downgraded for various reasons (risk of bias, inconsistency, indirectness, imprecision and publication bias). Consequently, the quality of the evidence was classified as high, moderate, low or very low. As described in table 2, we have for many of these outcomes downgraded for study limitations/high risk of bias or for imprecision because there were few events in many of these studies.

DISCUSSION

In this systematic review addressing the effect and safety of prehospital administration of ketamine in analgesic

doses, we included eight studies with 2760 patients in total.

Strengths and limitations of this systematic review

The included studies were heterogeneous in terms of setting, patient population and outcomes explored as well as in their comparators, such as intravenous or intranasal ketamine with a variety of opioids or with nitrous oxide. In addition, a single dose of ketamine was compared with ketamine that was administered continuously.

Although the evidence base includes five RCTs, five of the eight included studies have a high risk of bias. The RCTs were relatively small studies with 63, 65, 120, 135 and 308 patients included, respectively. None of the studies was designed or powered to truly test the safety of ketamine. Adverse events and the severity thereof were inconsistently reported.

The eight studies cover five different comparisons, so the amount of research evidence for each comparison is sparse. Only one of the outcomes in one of the comparisons has been measured in more than one study of similar design, and several of the outcomes have not been assessed in a prehospital study at all. When using GRADE to assess our confidence in the estimates, we more often than not, downgraded for high risk of bias or imprecision due to very few events or wide CIs. Three of the eight included studies are observational studies. They have an initial high risk of bias compared with RCTs due to the lack of randomisation. This is acknowledged in GRADE where observational studies start at low quality of evidence. Lack of blinding is a weakness in all of these studies. This becomes a challenge when the main outcome is subjective, pain, and we have downgraded for




Figure 8 Ketamine and N2O versus only N2O—adverse events.

Table 2 Summary of findings for the comparisons

Ketamine compared with opioids for prehospital pain management						
Patient or population: prehospital pain management setting: prehospital setting in the USA and Vietnam Intervention: ketamine Comparison: opioids						
Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with opioids	Risk with ketamine				
Change in pain score assessed with VAS	The mean change in the pain score was 3.1	The mean change in the pain score in the intervention group was 0.4 less (0.8 less to 0)	-	308 (1 RCT)	⊕⊕⊕○ Moderate †	
Change in pain score assessed with NRS scale from: 1 to 10	The mean change in the pain score was 2.5	The mean change in pain score in the intervention group was 3 less (3.86 less to 2.14 less)	-	158 (1 observational study)	⊕⊕○○ Low	
Serious adverse events	51 per 1000	0 per 1000 (0 to 0)	Not estimable	158 (1 observational study)	⊕⊕○○ Very low	
Nausea and vomiting	194 per 1000	47 per 1000 (21 to 101)	RR 0.24 (0.11 to 0.52)	308 (1 RCT)	⊕⊕⊕○ Moderate †	
Agitation	14 per 1000	112 per 1000 (27 to 474)	RR 7.81 (1.85 to 32.97)	308 (1 RCT)	⊕⊕⊕○ Moderate †	
Ketamine and morphine compared with only morphine for prehospital pain management						
Patient or population: prehospital pain management setting: prehospital setting in Sweden, France and Australia Intervention: ketamine and morphine Comparison: only morphine						
Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Certainty of the evidence (GRADE)	Comments
	Risk with only morphine	Risk with ketamine and morphine				
Change in pain scores Scale from: 1 to 10	The mean change in pain scores was 3.5	Mean 1.51 lower (3.36 lower to 0.33 higher)	-	135 (2 RCTs)	⊕⊕○○ Low ‡	
Change in pain scores Scale from: 1 to 10	The mean change in pain score was 3.1	Mean 1.3 lower (2.95 lower to 0.35 higher)	-	27 (1 observational study)	⊕○○○ Very low §	
Serious adverse events	Not reported	Not reported	Not estimable	-	-	None of the two studies reported any serious adverse events
Total number of adverse events	165 per 1 000	468 per 1 000 (289 to 764)	RR 2.84 (1.75 to 4.63)	200 (2 RCTs)	⊕⊕⊕○ Moderate ¶	
Continuous administration of ketamine compared with ketamine given as a bolus for prehospital pain management						

Continued

Table 2 Continued

Patient or population: prehospital pain management setting: prehospital setting in France
Intervention: continuous administration of ketamine
Comparison: ketamine given as a bolus

Outcomes	Anticipated absolute effects* (95% CI)	Risk with the continuous administration of ketamine	Relative effect (95% CI)	Number of participants (studies)	Certainty of the evidence (GRADE)	Comments
Change in pain scores. Scale from 0 to 10	Risk with ketamine given as a bolus The mean change in the pain score was 3.1	The mean change in pain score in the intervention group was 0.6 less (1.84 less to 0.64 more)	–	63 (1 RCT)	⊕⊕○○ Low**	
Serious adverse events	–	–	Not estimable	(One study)	–	No serious events were reported
Nausea and vomiting	91 per 1000	0 per 1000 (0 to 0)	Not estimable	63 (1 RCT)	⊕⊕○○ Very low††	

Ketamine and nitrous oxide compared with only nitrous oxide for prehospital pain management

Patient or population: prehospital pain management setting: prehospital setting in Canada
Intervention: ketamine and nitrous oxide
Comparison: only nitrous oxide

Outcomes	Anticipated absolute effects* (95% CI)	Risk with ketamine and nitrous oxide	Relative effect (95% CI)	Number of participants (studies)	Certainty of the evidence (GRADE)	Comments
≥2 point reduction in pain, 15 min	Risk with only nitrous oxide 350 per 1 000	634 per 1 000 (427 to 931)	RR 1.81 (1.22 to 2.66)	120 (1 RCT)	⊕⊕⊕○ Moderate††	
≥2 point reduction in pain, 30 min	407 per 1 000	758 per 1 000 (534 to 1 000)	RR 1.86 (1.31 to 2.66)	108 (1 RCT)	⊕⊕⊕○ Moderate ††	
Serious adverse events	0 per 1 000	0 per 1 000 (0 to 0)	Not estimable	(1 RCT)	–	
Total number of adverse events	233 per 1 000	866 per 1 000 (541 to 1 000)	RR 3.71 (2.32 to 5.31)	120 (1 RCT)	⊕⊕⊕○ Moderate††	
Number of patients with adverse events	200 per 1 000	616 per 1 000 (358 to 1 000)	RR 3.08 (1.79 to 5.31)	120 (1 RCT)	⊕⊕⊕○ Moderate††	

Continued

Table 2 Continued

*The risk in the intervention group (and its 95% CI) assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI). GRADE Working Group grades of evidence: high certainty: we are very confident that the true effect is similar to that of the estimated effect. Moderate certainty: we are moderately confident in the effect estimate; the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different. Low certainty: our confidence in the effect estimate is limited; the true effect may be substantially different from the estimated effect. Very low certainty: we have very little confidence in the effect estimate; the true effect is likely to be substantially different from the estimated effect.

†Downgraded one level for high risk of bias.

‡Downgraded one level for imprecision, Only four events and all four of them in the same group. There were no events in the other group and therefore RR cannot be estimated.

§Downgraded one level for imprecision, This cohort only has 27 patients included.

¶Downgraded one level for risk of bias due to unclear randomisation and open label design.

** Downgraded one level for imprecision, one study included with 63 patients.

††Downgraded one level for imprecision, Only three events.

‡‡Downgraded one level for imprecision, only one study with a total of 120 patients. There are also large effects, but with unclear blinding we do not upgrade. NRS, numeric pain rating scale; RCT, randomised controlled trial; VAS, visual analogue scale.

high risk of bias. However, there is moderate quality of the evidence for the main outcome, change in pain score, for one of the comparisons.

This systematic review has the benefit of systematic and transparent preplanned methodology. Decisions and judgements were conducted by two authors independently of each other, hence, reducing the risk of bias in the conduct of the review. We conducted a wide literature search in several databases, but it is still possible that there exist relevant studies that we did not identify, both in other databases and in other languages. As always with systematic reviews, there is the possibility that relevant studies may have been published after our search was conducted. Our deviation from the protocol to include a large study even though it included some children may be interpreted as a limitation. However, we would argue that the inclusion of extra patients (1876 patients added to the 884 patients from the other seven studies), where the large majority was adults adds greatly to the available information regarding side effects/adverse events.

Two of the studies were conducted in Iraq and in Vietnam, respectively, where a number of patients were injured in mine accidents. These studies were the largest studies and included 1909 patients. It is reasonable to assume that the results from studies conducted in war zones are not directly applicable in civilian settings since the victims tend to be men, relatively young and previously healthy and are not representative of trauma victims in general. The study from Iraq did not report on any of our predefined outcomes.

Clinical implications

Ketamine administered in analgesic doses (0.1–0.2 mg/kg) intravenously appears to be at least as effective as opioids administered alone considering pain reduction. In the study from Iraq, an initial dose of ketamine (0.2 mg/kg) was given in all cases of penetrating trauma and burns, but patients with TBI or blunt injury received only pentazocine.¹⁹ The patients in the study conducted by Tran *et al* received 0.2–0.3 mg/kg intravenous of ketamine.²⁶ Four^{22–25} of the other five studies included studies administered ketamine in 0.2–0.3 mg/kg intravenous doses, while in the last intravenous study,²⁰ the patients received 10–20 mg intravenous. of ketamine. In the study where ketamine was administered intranasally, the patients received an average of 0.75 mg/kg of ketamine.²¹ Hence, the patients in all studies received appropriate analgesic doses of ketamine.

Adverse events

In general, very few adverse events were reported in the included studies. Most of the events were related to nausea and vomiting. Agitation was more common in the ketamine group in the study performed by Tran *et al*.²⁶ Bronsky *et al* reported that two patients experienced respiratory compromise and two suffered haemodynamic instability.²⁵ All four patients were in the fentanyl group.

Given the safety profile of ketamine and the results reported in the included studies, it appears reasonable to suggest that low-dose ketamine for analgesic purposes can be administered safely during prehospital emergency care when proper indications and contraindications are identified. Prehospital healthcare providers with a level of training suitable to administer ketamine—that is, personnel that are trained to handle potential adverse events—must be identified. None of the included studies had enough power to detect differences in rare events, and the quality of evidence was poor. One of the studies showed an increased number (pooled) of adverse events in the group receiving ketamine and morphine, indicating that an improved analgesic effect increases the risk for adverse events. It is unclear whether adverse events are more likely to occur with opioids than with ketamine. However, it is essential to note that this review describes ketamine administered in analgesic doses and not in sedative and anaesthetic doses where advanced skills are required to be able to handle the patient in an adequate manner.

Studies from other settings

In a recent systematic review and meta-analysis, Yousefi-fard *et al* included seven studies and pooled the effect estimates of observational and randomised interventional studies.²⁹ They concluded that ketamine is an effective and safe medication in prehospital pain management in patients with trauma and can be considered as an acceptable alternative to opioids. The analgesic effect of low-dose ketamine is also employed in the hospital. In a recent systematic review and meta-analysis, Karlow *et al* studied ketamine as an alternative to opioids for acute pain in the emergency department (ED).³⁰ The authors concluded that ketamine can be used as an alternative to opioids in the ED, as they found that ketamine was non-inferior to opioids. They also found that the rate of non-severe adverse effects was higher with ketamine. It is unclear to what extent results from ED studies can be extrapolated to the prehospital setting. However, it is not obvious that the safety profile of ketamine in the prehospital setting is independent of the qualifications of the healthcare provider that administers the drug. Studies specifically addressing competence of prehospital providers administering ketamine should therefore be conducted. The body of evidence for benefit and possible harm is limited as few studies have been performed. Future studies need to address all relevant side effects, the optimal drug dose as well as all relevant outcome measures.

CONCLUSION

This systematic review of the current literature indicates that ketamine is an effective analgesic to be administered prehospitally.

Author affiliations

- ¹Division of Prehospital Services, Air Ambulance Department, Oslo University Hospital, Oslo, Norway
- ²Faculty of Medicine, University of Oslo, Oslo, Norway
- ³Faculty of Health Sciences, University of Stavanger, Stavanger, Norway
- ⁴Department of Research, Norwegian Air Ambulance Foundation, Oslo, Norway
- ⁵Trauma Unit, Sørlandet Hospital, Kristiansand, Norway
- ⁶Department of Prehospital Care and Disaster Medicine, Region of Skåne, Lund, Sweden
- ⁷Research Department, Prehospital Emergency Medical Service, Central Denmark Region, Aarhus, Denmark
- ⁸Department of Anaesthesiology, Regional Hospital of Horsens, Horsens, Denmark
- ⁹Centre for Prehospital Emergency Care, Oulu University Hospital, Oulu, Finland
- ¹⁰Anaesthesia Research group, MRC, Oulu University Hospital and University of Oulu, Oulu, Finland
- ¹¹Department of Anaesthesiology and Intensive Care, Linköping University Hospital, Linköping, Sweden
- ¹²Department of Biomedical and Clinical Sciences, Linköping University, Linköping, Sweden
- ¹³Landspítali University Hospital, Reykjavik, Iceland
- ¹⁴Danish Air Ambulance, Aarhus, Denmark
- ¹⁵Department of Anaesthesiology, Aarhus University Hospital, Aarhus, Denmark
- ¹⁶Department of Clinical Medicine, Aarhus University, Aarhus, Denmark
- ¹⁷Centre for Prehospital Emergency Medicine, Kuopio University Hospital, Kuopio, Finland
- ¹⁸University of Eastern Finland, Kuopio, Finland
- ¹⁹Division for Health Services, Norwegian Institute of Public Health, Oslo, Norway

Acknowledgements We thank research librarian Jane Kjemtrup Andersen (Viborg Regional Hospital, Denmark) for developing the search strategies.

Contributors MS supervised the process of drafting this manuscript and coordinated all identification, data extraction and appraisal of included manuscripts. MS and either PKH, MR or PK independently assessed all potential eligible articles for inclusion. GEV coordinated all methodological support. MS, PKH, PK, KDF, LR, RL, VM, LR, JK, MR, GEV are members of the SSAI task force on prehospital pain management and participated in planning, design, interpretation of results, manuscript drafting and revisions of the manuscript.

Funding SSAI funded meetings and software costs. The Norwegian Air Ambulance Foundation funded publication costs. Otherwise this research received no further grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Lasse Raatiniemi <http://orcid.org/0000-0001-5537-9749>
Leif Rognås <http://orcid.org/0000-0002-2542-565X>
Marius Rehn <http://orcid.org/0000-0001-9519-241X>

REFERENCES

- 1 Jennings PA, Cameron P, Bernard S. Epidemiology of prehospital pain: an opportunity for improvement. *Emerg Med J* 2011;28:530–1.
- 2 Albrecht E, Taffe P, Yersin B, *et al*. Undertreatment of acute pain (oligoanalgesia) and medical practice variation in prehospital analgesia of adult trauma patients: a 10 yr retrospective study. *Br J Anaesth* 2013;110:96–106.
- 3 Galinski M, Ruscev M, Gonzalez G, *et al*. Prevalence and management of acute pain in prehospital emergency medicine. *Prehosp Emerg Care* 2010;14:334–9.
- 4 Bounes V, Barniol C, Minville V, *et al*. Predictors of pain relief and adverse events in patients receiving opioids in a prehospital setting. *Am J Emerg Med* 2011;29:512–7.
- 5 Lodge D, Anis NA, Burton NR. Effects of optical isomers of ketamine on excitation of cat and rat spinal neurones by amino acids and acetylcholine. *Neurosci Lett* 1982;29:281–6.
- 6 Craven R. Ketamine. *Anaesthesia* 2007;62 Suppl 1:48–53.
- 7 Green SM, Johnson NE. Ketamine sedation for pediatric procedures: Part 2, review and implications. *Ann Emerg Med* 1990;19:1033–46.
- 8 Bourgoin A, Albanèse J, Léone M, *et al*. Effects of sufentanil or ketamine administered in target-controlled infusion on the cerebral hemodynamics of severely brain-injured patients. *Crit Care Med* 2005;33:1109–13.
- 9 Bourgoin A, Albanèse J, Wereszczynski N, *et al*. Safety of sedation with ketamine in severe head injury patients: comparison with sufentanil. *Crit Care Med* 2003;31:711–7.
- 10 Albanèse J, Arnaud S, Rey M, *et al*. Ketamine decreases intracranial pressure and electroencephalographic activity in traumatic brain injury patients during propofol sedation. *Anesthesiology* 1997;87:1328–34.
- 11 Bar-Joseph G, Guilburd Y, Tamir A, *et al*. Effectiveness of ketamine in decreasing intracranial pressure in children with intracranial hypertension. *J Neurosurg Pediatr* 2009;4:40–6.
- 12 Sherwin TS, Green SM, Khan A, *et al*. Does adjunctive midazolam reduce recovery agitation after ketamine sedation for pediatric procedures? A randomized, double-blind, placebo-controlled trial. *Ann Emerg Med* 2000;35:229–38.
- 13 McManus JG, Sallee DR. Pain management in the prehospital environment. *Emerg Med Clin North Am* 2005;23:415–31.
- 14 Kennedy RM, Porter FL, Miller JP, *et al*. Comparison of fentanyl/midazolam with ketamine/midazolam for pediatric orthopedic emergencies. *Pediatrics* 1998;102:956–63.
- 15 McGuinness SK, Wasiak J, Cleland H, *et al*. A systematic review of ketamine as an analgesic agent in adult burn injuries. *Pain Med* 2011;12:1551–8.
- 16 Bonanno FG. Ketamine in war/tropical surgery (a final tribute to the racemic mixture). *Injury* 2002;33:323–7.
- 17 Higgins JPT, Green S. *Cochrane handbook for systematic reviews of interventions*. The Cochrane Collaboration, 2011.
- 18 Atkins D, Best D, Briss PA, *et al*. Grading quality of evidence and strength of recommendations. *BMJ* 2004;328:1490–7.
- 19 Losvik OK, Murad MK, Skjerve E, *et al*. Ketamine for prehospital trauma analgesia in a low-resource rural trauma system: a retrospective comparative study of ketamine and opioid analgesia in a ten-year cohort in Iraq. *Scand J Trauma Resusc Emerg Med* 2015;23:94.
- 20 Jennings PA, Cameron P, Bernard S, *et al*. Morphine and ketamine is superior to morphine alone for out-of-hospital trauma analgesia: a randomized controlled trial. *Ann Emerg Med* 2012;59:497–503.
- 21 Andolfatto G, Innes K, Dick W, *et al*. Prehospital analgesia with intranasal ketamine (PAIN-K): a randomized double-blind trial in adults. *Ann Emerg Med* 2019;74:241–50.
- 22 Wiel E, Zitouni D, Assez N, *et al*. Continuous infusion of ketamine for out-of-hospital isolated orthopedic injuries secondary to trauma: a randomized controlled trial. *Prehosp Emerg Care* 2015;19:10–16.
- 23 Galinski M, Dolveck F, Combes X, *et al*. Management of severe acute pain in emergency settings: ketamine reduces morphine consumption. *Am J Emerg Med* 2007;25:385–90.
- 24 Johansson P, Kongstad P, Johansson A. The effect of combined treatment with morphine sulphate and low-dose ketamine in a prehospital setting. *Scand J Trauma Resusc Emerg Med* 2009;17:61.
- 25 Bronsky ES, Koola C, Orlando A, *et al*. Intravenous low-dose ketamine provides greater pain control compared to fentanyl in a civilian prehospital trauma system: a propensity matched analysis. *Prehosp Emerg Care* 2019;23:1–8.
- 26 Tran KP, Nguyen Q, Truong XN, *et al*. A comparison of ketamine and morphine analgesia in prehospital trauma care: a cluster randomized clinical trial in rural Quang tri Province, Vietnam. *Prehosp Emerg Care* 2014;18:257–64.
- 27 Champion HR, Sacco WJ, Copes WS, *et al*. A revision of the trauma score. *J Trauma* 1989;29:623–9.
- 28 Johansson J, Sjöberg J, Nordgren M, *et al*. Prehospital analgesia using nasal administration of S-ketamine—a case series. *Scand J Trauma Resusc Emerg Med* 2013;21:38.
- 29 Yousefifard M, Askarian-Amiri S, Rafiei Alavi SN, *et al*. The efficacy of ketamine administration in prehospital pain management of trauma patients; a systematic review and meta-analysis. *Arch Acad Emerg Med* 2020;8:e1.
- 30 Karlow N, Schlaepfer CH, Stoll CRT, *et al*. A systematic review and meta-analysis of ketamine as an alternative to opioids for acute pain in the emergency department. *Acad Emerg Med* 2018;25:1086–97.

SEARCH STRATEGIES

PUBMED

#1,"Search ketamine[Text Word]"
#2,"Search analgesics, ketamine[Pharmacological Action]"
#3,"Search ketamine[MeSH Terms]"
#4,"Search (#1 OR #2 OR #3)"
#5,"Search emergency medical services[MeSH Terms]"
#6,"Search Ambulances[MeSH Terms]"
#7,"Search Ambulance*[Text Word]"
#8,"Search Prehospital[Text Word]"
#9,"Search Pre-hospital[Text Word]"
#10,"Search out of hospital[Text Word]"
#11,"Search Paramed*[Text Word]"
#12,"Search emergency medical technicians[MeSH Terms]"
#13,"Search (#5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12)"
#14,"Search Danish[Language]"
#15,"Search Norwegian[Language]"
#16,"Search Swedish[Language]"
#17,"Search English[Language]"
#18,"Search (#14 OR #15 OR #16 OR #17)"
#19,"Search (#4 AND #13 AND #18)"
#20,"Search Animals[MeSH Terms]"
#21,"Search Humans[MeSH Terms]"
#22,"Search (#20 NOT #21)"
#23,"Search (#19 NOT #22)"
#24,"Search (""xxxx/xx/xx""[Date - Entrez]: ""xxxx/xx/xx""[Date - Entrez])"
#25,"Search (#23 AND #24)"

EMBASE

#1, analgesic agent/
#2, ketamine.m. _titl.
#3, rescue personnel/
#4, ambulance/
#5, emergency health service/
#6, "emergency medical technician*".ab.ti.
#7, "emergency responder*".ab.ti.
#8, rescue service. ab.ti.
#9, "Paramed*". ab.ti.
#10, "ambulance*". ab.ti.
#11, pre-hospital. ab.ti.
#12, prehospital. ab.ti.
#13, out-of-hospital. ab.ti.
#14, or/3-13
#15, or/1-2
#16, and/14-15

Cochrane Library

#1 MeSH descriptor: [Ketamine] explode all trees

#2 MeSH descriptor: [Emergency Medical Technicians] explode all trees

#3 MeSH descriptor: [Ambulances] explode all trees

#4 paramed*

#5 out-of-hospital

#6, pre-hospital

#7, prehospita

#8, ambulance*

#9, #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8

#10, #1 AND # 9

Epistemonikos

(title:((title:(prehospital OR pre-hospital OR out-of-hospital OR ambulance* OR “emergency medical technicians” OR “emergency medical service*”) OR abstract:(prehospital OR pre-hospital OR out-of-hospital OR ambulance* OR “emergency medical technicians” OR “emergency medical service*”))) AND (title:(ketamin*) OR abstract:(ketamin*)))

Appendix 2

Excluded reference	Reason for exclusion
Allison K, Porter K. Consensus on the pre-hospital approach to burns patient management. <i>Injury</i> 2004; 35: 734-8.	Ketamine not central in the text
Anonymous. Comments on the recommendations of the German Medical Society dated October 20, 2003 on the administration of analgesics by paramedics in emergency situations. <i>Notarzt</i> 2005; 21: 81-82.	Excluded due to language restrictions
Ansem RP, Hartman JA, Foudraine JF, van Loenen E, Rutten FL. [Analgetic ketamine feasible in ambulance emergency care]. <i>Ned Tijdschr Geneeskd</i> 1994; 138: 2301-4.	Excluded due to language restrictions
Ardeel E. Adverse effects following prehospital use of ketamine by paramedics. <i>Acad Emerg Med</i> 2012; 19: S269-S70.	Letter
Aries P, Montelescaut E, Pessey F, Danguy des Deserts M, Giacardi C. Pre-hospital emergency medicine: pain control. <i>Lancet</i> 2016; 387: 747.	Ketamine not central in the text
Arroyo-Novoa CM, Figueroa-Ramos MI, Miaskowski C, Padilla G, Paul SM, Rodriguez-Ortiz P, Stotts NA, Puntillo KA. Efficacy of small doses of ketamine with morphine to decrease procedural pain responses during open wound care. <i>Clin J Pain</i> 2011; 27: 561-6.	Patient population did not match the criteria
Barrett TW, Schriger DL. Move over morphine: Is ketamine an effective and safe alternative for treating acute pain? Answers to the September 2015 journal club. <i>Ann Emerg Med</i> 2016; 67: 289-94.	Setting did not match the criteria
Berg C. Out-of-hospital ketamine for pain, agitation, and airway intervention is safe and effective. <i>Ann Emerg Med</i> 2015; 66: S32.	Study design did not match the criteria
Bredmose PP, Grier G, Davies GE, Lockey DJ. Pre-hospital use of ketamine in paediatric trauma. <i>Acta Anaesthesiol Scand</i> 2009; 53: 543-5.	Patient population did not match the criteria

Bredmose PP, Lockey DJ, Grier G, Watts B, Davies G. Pre-hospital use of ketamine for analgesia and procedural sedation. EMJ 2009; 26: 62-4.	Study design did not match the criteria
Brokmann JC, Rossaint R, Hirsch F, Beckers SK, Czaplik M, Chowanetz M, Tamm M, Bergrath S. Analgesia by telemedically supported paramedics compared with physician-administered analgesia: A prospective, interventional, multicentre trial. Eur J Pain 2016; 20: 1176-84.	Ketamine not central in the text
Butler FK, Kotwal RS, Buckenmaier CC, 3rd, Edgar EP, O'Connor KC, Montgomery HR, Shackelford SA, Gandy JV, 3rd, Wedmore IS, Timby JW, Gross KR, Bailey JA. A triple-option analgesia plan for tactical combat casualty care: TCCC guidelines change 13-04. J Spec Oper Med 2014; 14: 13-25.	Study design did not match the criteria
Castle N, Naidoo R. Achieving prehospital analgesia. EMJ 2012; 29: 765-6.	Setting did not match the criteria
Castren M, Lindstrom V, Branzell JH, Niemi-Murola L. Prehospital personnel's attitudes to pain management. Scand J Pain 2015; 8: 17-22.	Study design did not match the criteria
Chesters A, Webb T. Ketamine for procedural sedation by a doctor-paramedic prehospital care team: a 4-year description of practice. Eur J Emerg Med 2015; 22: 401-6.	Patient population did not match the criteria
Corrigan M, Wilson SS, Hampton J. Safety and efficacy of intranasally administered medications in the emergency department and prehospital settings. Am J Health Syst Pharm 2015; 72: 1544-54.	Study design did not match the criteria
Domonoske B, Gunter R, Love J. Ketamine may increase the risk of PE in selected trauma patients. Crit Care Med 2014; 42: A1610.	Setting did not match the criteria
Domonoske B, Love J. Ketamine reduces the incidence of VTE in selected trauma patients. Crit Care Med 2013; 41: A55.	Setting did not match the criteria
Eidenbenz D, Taffe P, Hugli O, Albrecht E, Pasquier M. A two-year retrospective review of the determinants of pre-hospital analgesia administration by alpine helicopter	Study design did not match the criteria

emergency medical physicians to patients with isolated limb injury. <i>Anaesthesia</i> 2016; 71: 779-87.	
Ellerton J, Paal P, Brugger H. Prehospital use of ketamine in mountain rescue. <i>EMJ</i> 2009; 26: 760-1.	Letter
Fisher AD, Rippee B, Shehan H, Conklin C, Mabry RL. Prehospital analgesia with ketamine for combat wounds: a case series. <i>J Spec Oper Med</i> 2014; 14: 11-7.	Study design did not match the criteria
Galinski M, Hoffman L, Bregeaud D, Kamboua M, Ageron FX, Rouanet C, Hubert JC, Istria J, Ruscev M, Tazarourte K, Pevirieri F, Lapostolle F, Adnet F. Procedural sedation and analgesia in trauma patients in an out-of-hospital emergency setting: A prospective multicenter observational study. <i>Prehosp Emerg Care</i> 2018; 22: 497-505.	Patient population did not match the criteria
Gausche-Hill M, Brown KM, Oliver ZJ, Sasson C, Dayan PS, Eschmann NM, Weik TS, Lawner BJ, Sahni R, Falck-Ytter Y, Wright JL, Todd K, Lang ES. An evidence-based guideline for prehospital analgesia in trauma. <i>Prehosp Emerg Care</i> ; 18 25-34.	Study design did not match the criteria
Green SM, Roback MG, Krauss B, Brown L, McGlone RG, Agrawal D, McKee M, Weiss M, Pitetti RD, Hostetler MA, Wathen JE, Treston G, Garcia Pena BM, Gerber AC, Losek JD. Predictors of emesis and recovery agitation with emergency department ketamine sedation: an individual-patient data meta-analysis of 8,282 children. <i>Ann Emerg Med</i> 2009; 54: 171-80.	Patient population did not match the criteria
Guldner GT, Petinaux B, Clemens P, Foster S, Antoine S. Ketamine for procedural sedation and analgesia by nonanesthesiologists in the field: a review for military health care providers. <i>Mil Med</i> 2006; 171: 484-90.	Patient population did not match the criteria
Gurnani A, Sharma PK, Rautela RS, Bhattacharya A Analgesia for acute musculoskeletal trauma: low-dose subcutaneous infusion of ketamine. <i>Anaesth Intensive Care</i> 1996; 24: 32-6	Patient population did not match the criteria
Henderson L. Special K for special situations. A review of ketamine for prehospital use. <i>JEMS</i> 2016; 41: 58-60.	Study design did not match the criteria

Hossfeld B, Holstrater S, Bernhard M, Lampl L, Helm M, Kulla M. Prehospital analgesia in adults. <i>Anesthesiol Intensivmed Notfallmed Schmerzther</i> 2016; 51: 84-96.	Excluded due to language restrictions
Iqbal M, Spaight PA, Siriwardena AN. Patients' and emergency clinicians' perceptions of improving pre-hospital pain management: A qualitative study. <i>EMJ</i> 2013; 30: e18.	Ketamine not central in the text
Jansen A, Boyle M. Prehospital pain relief, where are we now? A review of the literature. <i>Australas J Paramed</i> 2014; 11: 20.	Ketamine not central in the text
Jennings PA, Cameron P, Bernard S. Ketamine as an analgesic in the pre-hospital setting: a systematic review. <i>Acta Anaesthesiol Scand</i> 2011; 55: 638-43.	Study design did not match the criteria
Jennings PA, Cameron P, Bernard S. Determinants of clinically important pain severity reduction in the prehospital setting. <i>EMJ</i> 2012; 29: 333-34.	Ketamine not central in the text
Jennings PA, Cameron P, Bernard S, Walker T, Jolley D, Fitzgerald M, Masci K. Long-term pain prevalence and health-related quality of life outcomes for patients enrolled in a ketamine versus morphine for prehospital traumatic pain randomised controlled trial. <i>EMJ</i> 2014; 31: 840-43.	Setting did not match the criteria
Jennings PA, Cameron P, Bernard SA, Walker T, Fitzgerald M, Masci K. Ketamine is superior to morphine alone for the management of traumatic pain in the prehospital setting: A randomised controlled trial. <i>Emerg Med Australas</i> 2012; 24: 19.	Letter
Johansson J, Sjöberg J, Nordgren M, Sandstrom E, Sjöberg F, Zetterstrom H. Prehospital analgesia using nasal administration of S-ketamine--a case series. <i>Scand J Trauma Resusc Emerg Med</i> 2013; 21: 38.	Study design did not match the criteria
Kovar JL, Gleisberg GR, Ardeel ER, Basnawi A, Escott MEA. Hemodynamic changes in patients receiving ketamine sedation by emergency medical services. <i>Acad Emerg Med</i> 2012; 19: S87.	Study design did not match the criteria

Lovrinčević M, Kotob F, Santarosa J. Pain management in the trauma setting. <i>Semin Anesth Perioper Med Pain</i> 2005; 24: 34-40.	Setting did not match the criteria
Madeira F, Ferreira P, Lapa T, Tavares E. Prehospital pain management: Do we have to learn more about it? <i>Eur J Anaesthesiol</i> 2013; 30: 203-04.	Ketamine not central in the text
Marland S, Ellerton J, Andolfatto G, Strapazzon G, Thomassen O, Brandner B, Weatherall A, Paal P. Ketamine: Use in anesthesia. <i>CNS Neurosci Ther</i> 2013; 19: 381-89.	Setting did not match the criteria
McKay WP. Intravenous analgesia for out-of-hospital traumatic pain in adults: ketamine gives a greater reduction in pain than morphine but causes more adverse effects. <i>Evid Based Nurs</i> 2013; 16: 58-9.	Letter
McQueen C, Crombie N, Cormack S, Wheaton S. Prehospital use of ketamine for analgesia and procedural sedation by critical care paramedics in the UK: A note of caution? <i>EMJ</i> 2014; 31: 1029.	Letter
Moy R, Wright C. Ketamine for military prehospital analgesia and sedation in combat casualties. <i>J R Army Med Corps</i> 2018; 164: 436-37.	Study design did not match the criteria
Moy RJ, Le Clerc S. Ketamine in prehospital analgesia and anaesthesia. <i>Trends Anaesth Crit Care</i> 2011; 1: 243-45.	Study design did not match the criteria
Petz LN, Tyner S, Barnard E, Ervin A, Mora A, Clifford J, Fowler M, Bebartá VS. Prehospital and en route analgesic use in the combat setting: a prospectively designed, multicenter, observational study. <i>Mil Med</i> 2015; 180: 14-18.	Study design did not match the criteria
Porter K. Ketamine in prehospital care. <i>EMJ</i> 2004; 21: 351-4.	Study design did not match the criteria
Radvansky BM, Puri S, Sifonios AN, Eloy JD, Le V. Ketamine-a narrative review of its uses in medicine. <i>Am J Ther</i> 2016; 23: e1414-e26.	Study design did not match the criteria
Schauer SG, Mora AG, Maddry JK, Bebartá VS. Multicenter, prospective study of prehospital administration of analgesia in	Study design did not match the criteria

the U.S. combat theater of Afghanistan. <i>Prehosp Emerg Care</i> 2017; 21: 744-49.	
Schonenberg M, Reichwald U, Domes G, Badke A, Hautzinger M. Effects of peritraumatic ketamine medication on early and sustained posttraumatic stress symptoms in moderately injured accident victims. <i>Psychopharmacology (Berl)</i> 2005; 182: 420-5.	Study design did not match the criteria
Scott S, Paul B. UK and Victorian, Acute pain guidelines compared. <i>Australas J Paramed</i> 2013; 10: 32.	Letter
Steel A, Wharton R, Bates A, French J, Lewis S, Mackenzie R. Ketamine use in prehospital critical care. <i>EMJ</i> 2008; 25: 618-19.	Letter
Svenson JE, Abernathy MK. Ketamine for prehospital use: new look at an old drug. <i>Am J Emerg Med</i> 2007; 25: 977-80.	Patient population did not match the criteria
Wedmore IS, Butler FK. Battlefield analgesia in tactical combat casualty care. <i>Wilderness Environ Med</i> 2017; 28: S109-S16.	Study design did not match the criteria
Wood PR. Ketamine: Prehospital and in-hospital use. <i>Trauma</i> 2003; 5: 137-40.	Study design did not match the criteria
Zhang M, Cowan T, Smiles JP, Morgan M, Armstrong J, Goswami C, Sewell C. Prehospital analgesic choice in injured patients does not impact on rates of vomiting: Experience from a New South Wales primary retrieval service. <i>Emerg Med Australas</i> 2017; 30: 406-11.	Study design did not match the criteria